

Effect of Supplementary Irrigation Schedules and Bio-Fertilization on Yield and Yield Attributes of Faba Bean (*Vicia faba* L.), and Lentil (*Lens culinaris* L.), under Rainfed Conditions

Attia M. A.

Dept. of Plant Production, Desert Research Center, El-Matarya, Cairo, Egypt.

Received on: 6/3/2013

Accepted: 29/4/2013

ABSTRACT

Two field experiments were carried out during the two successive growing seasons (2006/2007 and 2007/2008) at El-Hammam farm, Marsa Matrouh, North Western Coast of Egypt, to study the effect of supplementary irrigation schedule and bio-fertilization on Faba bean (*Vicia faba* L.) and Lentil (*lens culinaris*, Medic) yields and its attributes.

The study included two regimes of bio-fertilizers (without bio-fertilization and bio-fertilization) with combined bio-fertilizer (*Rhizobium leguminosarum* bv. *viciae* and *Bacillus megatherium* var. *phosphaticum* strains) and three irrigation schedule (without supplementary irrigation (rain fed), one supplementary irrigation 45 mm and two supplementary irrigation 90 mm). The obtained results showed that:

- adding two supplementary irrigations(90mm) caused the highest values for all measured characteristics of yield and its components in both studied crops.
- The measurements of yield and its components i.e., plant height at harvest (cm), number of branches/plant; number of branches/m²; number of grains/plant; number of grains/pod; 100 grains weight (g); and biological, straw and grains yields (kg/fed) were significantly increased by adding bio-fertilizer than the control.
- In respect to the interaction between the two studied factors, biofertilizer and two supplementary irrigation with 90 mm/fed gave the highest values for all measured characteristics in both crops,

Key words: Faba bean (*Vicia faba* L.), Lentil (*lens culinaris* L.), bio-fertilization, rainfall, supplementary irrigation.

INTRODUCTION

For over half a century, the world has relied on increasing crop yields to supply an ever increasing demand for food. One of the major problems that limit economically successful agricultural production worldwide is poor soil fertility. Nutrient depletion of soil is a particular problem for small land holders in developing countries, where much grain-legume production occurs (Baset and Shamsuddin, 2010). It was suggested that the average annual nutrient depletion rates across 37 African countries were 22 kg N /ha, 25kg P/ha and 15kg K/ha (Sanchez, 2002). Therefore, addition of fertilizers is necessary to correct poor soil fertility by supplying nutrients needed for optimum crop growth (ELsheikh et al., 2005).

In the last three decades, the use of nitrogen fertilizer has increased almost nine folds (Vance, 2001). The increasing use of nitrogen and phosphorus fertilizers in high productive systems have created environmental problems such as deterioration of soil quality, surface and groundwater as well as air pollution, reduced biodiversity and suppressed ecosystem function (Schultz et al, 1998, Socalow, 1999 and Vance, 2001).

Crop Scientists all over the world are facing the alarming situation induced by the intensive uses of chemical fertilizers and they are trying to overcome this condition by exploring alternative source which is cost effective and save the environment. Biofertilizers, are alternate source of chemical fertilizer (Baset and Shamsuddin, 2010).

Legumes are the major source of proteins for both man and livestock, especially in poor countries, where animal protein is expensive (Hubbell and Gerald, 2003). It is the main food for millions of people and the source of protein for the middle and low income groups (Salih, 1981).

Grain legumes play an essential role in human nutrition balancing the deficiencies of cereal-based diet (Dart and Krantz, 1977). The importance of legumes is that they can fix nitrogen in symbiotic association with rhizobia, and so they increase the soil nitrogen content (Poth et al., 1986). This association enables legumes to benefit from an augmented nitrogen supply and can grow well on relatively poor soils (Heywood et al., 1985).

Faba bean (*Vicia faba* L.) and Lentil (*Lens culinaris*, Medic) are grown world-wide in cropping systems as a grain (pulse) and green-manure legume. The grain legumes contribute to the sustainability of cropping systems via: 1) its ability

to contribute nitrogen (N) to the system via biological N fixation, 2) diversification of systems leading to decreased disease, pest and weed build-up and potentially increased biodiversity, 3) reduced fossil energy consumption in plant production, and 4) providing food and feed rich in protein.

Grain legumes are major crops cultivated in the Northern and the River Nile Governorates of Egypt. For example, Faba bean (*Vicia faba* L.) is produced in an average area of 69720 ha, with an average yield of 1896 kg/ha (AOAD, 2007). Grain legumes are important cash crops for farmers (Watson, 1981) under dry lands conditions. Supplementary irrigation could significantly prevent from undesirable effects of water deficit. Oweis et al. (2004) reported seed and biological yield in lentil increased by supplementary irrigation. Increasing seed and biological yield through supplementary irrigation reported by many researchers Zhang, H. et al. (2000). Therefore, the objectives of this study were to determine the effects of biological fertilization and supplementary watering on faba bean and lentil productivity under rainfed conditions at the North Western Coast of Egypt.

MATERIALS AND METHODS

Site description

Two field experiments were conducted at El-Hammam farm, Matrouh Governorate, North Western Coast of Egypt for two consecutive winter seasons 2006/2007 and 2007/2008 to study the effect of supplementary irrigation schedule), and bio-fertilizer application on faba bean and lentil crop productivity.

Meteorological data of temperature, dew point and relative humidity which were obtained from Egyptian Meteorological Authority, Agricultural Research Center during the two growing seasons 2006-2007 -2008 are shown in table (1).

Treatments

The experiments included 6 treatments, which were the combination of three irrigation schedules as supplementary irrigation treatments and two bio-fertilizers treatments as follows:

Irrigation schedule:

- 1-without supplementary irrigation (rainfall only), control (W0).
- 2-one supplementary irrigation by 45 mm (189m³/fed) after 45 days from sowing (W1).
- 3-two supplementary irrigations by 45 mm (378m³/fed) of each, the first after 45 days from sowing and the second at flowering stage (W2).

All the experimental plots received irrigation by 45 mm immediately after sowing to obtain good plant establishment.

Bio-fertilizer application:

- 1-without bio-fertilization (control) (F0).

- 2- inoculation with combined bio-fertilizer (*Rhizobium leguminosarum* bv. *viciae* and *Bacillus megatherium* var. *phosphaticum* strains) (F1).

Experimental design and execution

A separate experiment was devoted to each investigated legume crop, where the treatments were arranged in split plot design in four replications, the main plots were occupied with supplemental irrigation treatments and the subplots were devoted to bio-fertilizer treatments.

The land was prepared by deep poughing, harrowing, leveling and ridging. The land was then divided into plots 6 x 3.8 m each., 50 kg N/fed as urea was broadcasted immediately after sowing. Phosphorus was applied at a rate of 45 kg P₂O₅/fed as triple super phosphate broadcasted before sowing.

Certified seeds of faba bean (*Vicia faba* L.) cultivar (Giza 5) and Lentil (*Lens culinaris* Medic.) cultivar (Giza 9) were obtained from Legumes Research Division, Agriculture Research Center, Ministry of Agriculture and Reclamation Giza, Egypt.

Seeds prepared was carried by soaking in tap water for 24 hours and air dried for one hour then , the seeds were divided into two equal parts, the first one was without bio-fertilizer (control), and the second was inoculated with the combined bio-fertilizer. Two to three seeds were placed in a hole on the top of the ridge with 20 cm spacing (between holes) and 70 cm (between ridges).

The supplementary irrigation treatments were added by gated pipe distribution system and the irrigation water quantities measured through flow meter. irrigation system. The soil was fallow in the summer before sowing each of faba bean and lentil in both seasons. Seeds were sown in 15th November 2006 and 25th November 2007 at rate of 30 kg/fed, in the first and second seasons, respectively. While The harvesting dates were 1st April 2006 and 8th April 2008 In the first and second seasons, respectively.

Yield measurements

At maturity, each plot was harvested separately by cutting the plants. Plants were then threshed on a large mat, then collected to measure yield and its components of each plot.

The following parameters, of each of faba bean and lentil yield and its components, were recorded:

1. plant height at harvest (cm),
2. number of branches/plant;
3. number of branches/m²;
4. number of pods/plant;
5. number of grains/pod;
6. 100 grains weight (g); and
7. biological, straw and grains yields (kg/fed)

Table 1: Meteorological parameters of the 2006-2007 and 2007 -2008 seasons

Month	Temp. (°c)			Dew point (mm)			Humidity(%)	
	Min.	Max.	Avg.	Min.	Max.	Avg	Min.	Max.
2006/07								
Nov.	16.7	20.0	18.40	8.3	11.2	9.75	50.3	63.7
Dec.	12.3	15.6	13.95	5.0	10.8	7.9	54.7	67.7
Jan.	10.5	18.2	14.35	4.1	10.4	7.25	59.7	69.7
Feb.	10.6	18.5	14.55	5.1	11.7	8.4	57.0	69.7
March.	12.0	20.9	16.45	5.9	12.4	9.15	58.7	67.0
April	13.7	20.8	17.25	9.0	12.9	10.95	49.0	68.7
May.	17.6	26.0	21.8	7.3	17.3	12.3	59.3	68.7
2007/08								
Nov.	13.9	24.6	19.25	7.2	15.6	11.4	55.0	63.0
Dec.	10.7	20.2	15.45	6.3	11.1	8.7	53.3	64.0
Jan.	10.1	17.1	13.6	7.2	9.9	8.55	63.7	78.2
Feb.	9.6	17.2	13.4	6.1	8.9	7.5	61.9	76.2
March.	12.4	23.0	17.7	5.5	11.4	8.45	57.7	75.2
April	14.0	24.3	19.15	7.9	12.4	10.15	53.0	70.4
May.	16.2	25.5	20.85	11.9	15.7	13.8	56.4	71.7

Source: Egyptian Meteorological Authority, Cairo, Egypt.

The soil texture (0-20cm) is loamy sand . it's parent material varies between calcium carbonate and gypsum with 34.8-41.2 caco3 %

. Statistical analysis

Multifactor Analysis of Variance (ANOVA) was used to determine the effect of different treatments on measured parameters. Least significance difference was used to compare between means (Gomez and Gomez, 1984). Data combined analysis of the two experiments was done according to Lazic (2008). Significance was accepted at $p < 0.05$.

RESULTS AND DISCUSSION

. Effect of supplementary irrigation:

Table (2) showed that all measured parameters i.e., plant height at harvest (cm), number of branches/plant; number of branches/m²; number of pods/plant; number of grains/pod; 100 grains weight (g); and biological, straw and grains yields (kg/fed) were significantly increased by supplementary irrigation schedule 45 and 90 mm viz, non irrigation (rain), in both seasons.

The highest values of mentioned characters were obtained by adding 90 mm as a supplementary irrigation.

The superiority of supplemental irrigation at rate of 90 mm on the yield and its components in both seasons may be due to the increment in number of pods/m², number of grains/m², and weight of 100 grains as indicated in Table (2).

These findings agree with those obtained by Dart and Krantz, (1977) and Ahmed et. al., (2007) Increasing growth characters as a result of adding water supplemental irrigation at a rate of 90 mm caused superiority in yield and its components. These results may be due to the effect of water on encouraging cell division, elongation and turgidity which in turn increase dry matter.

In this respect, it is worthy mention that the major problem facing grain legumes production in Egypt northern governorates is synchronizing of inadequate rainfall incidences during the most drought susceptible stage of growth and development (Abdel, 1993). Most studies on grain legumes for improving yield confirmed that pod development and seed filling stages were the most drought sensitive (Abdel, 1982 and El-Hamadany, 2005).

. Effect of bio-fertilization:

Data in Table (3) showed that the yield and its components was increased significantly in all characters with application of bio-fertilizer i.e., plant height at harvest (cm), number of branches/plant; number of branches/m²; number of grains/plant; number of grains/pod; 100 grains weight (g); and biological, straw and grains yields (kg/fed) than those of the control (without bio-fertilizer). There were significant differences in all yield characters and these differences were fairly true under the probability of 5% level of significance. Bio-fertilizers had markedly affected all mentioned characters compared to those plants without bio-fertilizer. The increment in yield and its components of faba bean and lentil may be in directly due the effects of bio-fertilization package on enhancing vegetative growth, root growth and dry matter accumulation.

Generally, under bio-fertilizer treatment, grain, straw and biological yields, were increased in both seasons as shown in Table (3), and the bio-fertilized Faba bean Giza 5, scored the highest values of all characters yield and its components. These results are in harmony with those reported by many

Table 2: Supplementary irrigation effect on faba bean and lentil yield and its components (means of two seasons)

Treatment	Characters									
	Plant height (cm)	Number of pods/plant	Number of branches/plant	Number of branches/m ²	100 grains weight (g)	Number of grains/pod	Grains yield (kg/fed)	Straw yield (kg/fed)	Biological yield (kg/fed)	
	Faba bean									
W0	36.50	6.31	1.53	35.60	63.60	2.16	519.20	1056.90	1576.20	
W1	46.10	7.51	2.16	36.60	73.70	2.69	682.90	1543.30	2226.30	
W2	56.20	9.57	3.12	44.90	83.10	3.53	977.50	2376.80	3315.60	
LSD (5%)	2.95	0.38	0.16	10.01	3.09	0.26	68.80	159.70	174.30	
	Lentil									
W0	26.50	16.63	2.76	41.40	3.18	4.75	173.07	398.05	573.12	
W1	36.10	17.83	3.39	50.85	3.69	5.92	227.63	523.56	753.19	
W2	46.20	19.89	4.35	65.25	4.16	7.77	325.83	749.42	1077.25	
LSD (5%)	4.58	0.92	0.26	8.32	0.12	0.57	22.93	52.75	77.68	

W0 = rainfall only W1 = rainfall + one supplementary irrigation

W2 = rainfall + two supplementary irrigations

Table 3: Bio-fertilizer effect on faba bean and lentil yield and its components (means of two seasons)

Treatment	Characters									
	Plant height (cm)	Number of pods/plant	Number of branches/plant	Number of branches/m ²	100 grains weight (g)	Number of grains/pod	Grains yield (kg/fed)	Straw yield (kg/fed)	Biological yield (kg/fed)	
	Faba bean									
F0	38.6	6.42	1.78	71.3	65.6	2.33	560.7	1251.7	1828.1	
F1	43.2	7.39	2.22	38.3	70.3	2.68	684.8	1536.0	2182.6	
LSD (5%)	2.2	0.27	0.16	7.11	1.46	0.12	42.9	80.8	140.7	
	Lentil									
F0	28.60	16.74	3.01	45.15	3.28	5.13	186.90	429.87	618.77	
F1	33.20	17.71	3.45	51.75	3.52	5.90	228.27	525.01	755.28	
LSD (5%)	1.62	0.53	0.39	4.68	ns	0.26	14.30	32.89	49.19	

F0 = without bio-fertilization F1 = inoculation with combined bio-fertilizer

Table 4: Effect of the interaction between supplementary irrigation and bio-fertilizer on faba bean and lentil yield and its components (means of two seasons)

Treatment	Characters							Biological yield (kg/fed)	
	Plant height (cm)	Number of pods/plant	Number of branches/plant	Number of branches/m ²	100 grains weight (g)	Number of grains/pod	Grains yield (kg/fed)		Straw yield (kg/fed)
Faba bean									
W0 + F0	34.9	5.90	1.36	27.0	61.2	1.95	469.4	938.6	1407.9
W0 + F1	37.9	6.72	1.69	44.3	65.9	2.38	569.1	1175.4	1744.5
W1 + F0	43.3	7.20	1.97	35.0	71.3	2.56	631.1	1410.7	2069.7
W1 + F1	48.8	7.82	2.36	38.3	76.1	2.83	734.9	1675.9	2382.8
W2 + F0	64.2	9.02	2.70	42.8	81.1	3.38	870.4	2149.9	3067.9
W2 + F1	58.3	10.12	3.55	47.0	85.0	3.68	1084.7	2603.6	3563.2
LSD (5%)	4.44	0.55	0.32	14.2	2.92	0.24	85.8	161.6	281.4
Lentil									
W0 + F0	24.90	16.12	2.54	36.83	2.91	3.90	161.86	258.98	424.04
W0 + F1	27.90	16.94	2.87	41.62	3.14	4.76	196.24	313.99	513.43
W1 + F0	33.30	17.42	3.15	45.68	3.40	5.12	217.62	348.19	569.01
W1 + F1	38.80	18.04	3.54	51.33	3.62	5.66	253.41	405.46	662.08
W2 + F0	54.20	19.24	3.88	56.26	3.86	6.76	300.14	480.22	783.56
W2 + F1	48.30	20.34	4.73	68.59	4.05	7.36	374.03	598.46	975.69
LSD (5%)	3.58	0.61	0.47	8.43	0.18	0.47	28.63	52.38	85.29

W0 = rainfall only W1 = rainfall + one supplementary irrigation

W2 = rainfall + two supplementary irrigations

F0 = without bio-fertilization F1 = inoculation with combined bio-fertilizer

researchers (Beck and Duc, 1991, Anderson and Ingram, 1993 and ELsheikh et al., 2005).

It is reported that sink capacity of the plant is dependent mainly on vegetative growth of plant, vigorous vegetative growth which increased due to the application of bio-fertilizer and supply of photosynthesis for formation of more branches and pods of legumes (Baset and Shamsuddin, 2010).

. Effect of the interaction between supplementary irrigation schedule and bio-fertilizer under rainfed conditions.

Data in Table (4) showed that the effect of the interaction between bio-fertilizer and supplementary irrigation schedule it had significant effect on all studied characters of faba bean and lentil except for number of branches/m² which tended to behave in different way. The highest values of yield and its components were obtained by bio-fertilizing each of faba bean and lentil plants and adding supplemental irrigation water by 90mm. However, the lowest values for the mentioned characters were obtained without bio-fertilizer (control) and without supplementary irrigation (rainfed). These results are in agreement with those reported by (Osman et al., 2010).

CONCLUSION

The areas of the North Western Coast of Egypt are characterized by low rainfall rate (average of the last 50 years is 133 mm/year), but evapotranspiration is low due to mild temperatures, cloudy skies, formation of dew on the plants during he growing seasons, and soil texture which varies between sandy and sandy loamy that has reasonable water holding capacity, hence the decision makers planed to install some great canals to convey the Nile water to the different parts of Egyptian desert, such as El-Hammam canal, which was designed to support the rainfed agricultural with supplementary irrigation to get improved yields. The most important crops in this areas are grain legumes.

Under rainfed conditions in El-Hammam farm at North Western Coast of Egypt, it can be concluded that cultivating the bio-fertilized legumes and supplementary irrigation with 90 mm is recommended to obtain the highest yield for each of faba bean and lentil plants under rain fed conditions.

REFERENCES

- Abdel, C. G. (1982). Drought resistance in (*Vicia faba* L.): A study of four cultivars. M.sc Thesis, Bath University, England, UK.
- Abdel, C. G. (1993). Effect of complementary watering on growth stages and yield of (*Vicia faba* L.). Mesopotamia J. of Agric. Vol (25), No 3, pp 5-10.
- Ahmed, E.A., A. Mohamed and K.H. Siddig, (2007). Economics of faba bean production and marketing in Northern Sudan. Conference on International Agricultural Research for Development. University of Kassel-Witzenhausen and University of Gottingen.
- Anderson, J.M. and J.S. Ingram, (1993). Tropical Soil Biology and Fertility. A Handbook of Methods. 2nd edition. CAB International. Wallingford. UK.
- AOAD, (2007). Arab Agriculture Statistics Year Book, 27 .Arab Organization for Agricultural Development .Khartoum, Sudan .
- Baset Mia, M.A. and Z.H. Shamsuddin, (2010). Rhizobium as a crop enhancer and biofertilizer for increased cereal production. African journal of Biotechnology, 9(37):6001-6009.
- Beck, D. and G. Duc, (1991). Improving N₂ fixation in faba bean. In: Rhizobium Inoculation and N Nutrition Options. Mediterraneanes-Serie Seminaires -no 10: 97-103.
- Dart, P.J. and B.A. Krantz, (1977). Legumes in the semiarid tropics. In: Exploiting the legume - Rhizobium symbiosis in tropical agriculture. (Eds. Vincent, J.M., Whitney, A.S. and Bose, J.). University of Hawaii, NifTAL project, USA., pp: 119-154 .
- El-Hamadany, S. Y. (2005). The effects of supplemental Irrigation and Abscisic acid (ABA) spraying on growth and yield of some faba bean (*Vicia faba* L.) cultivars. Ph.D Thesis , Mosul University, Mosul, Iraq.
- ELsheikh, M.A., A.M.A. EL-Tilib and E.A.E. ELsheikh, (2005). A Note on the effect of phosphate rock, triple superphosphate, Bradyrhizobium and their combination on the available soil phosphorus in shambat clay soil. University of Khartoum Journal of Agricultural Sciences, 13(3): 488-493.
- Gomez, K.A. and A.A. Gomez (1984). Statistical procedures for agriculture Research. A willey-Inter Sceince Publication. John Willey & Sons Inc., New York, USA.
- Heywood, V.H., D.M. Moore, L.B.K. Richardson and W.T . Stearn (1985). Flowering plants of the world . Croom Helm, London and Sydney, pp: 122-144 .
- Hubbell, D.H. and K. Gerald, (2003). Biological nitrogen fixation. Fact sheet of the Soil and Water Science Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Science, University of Florida, pp: 4 .
- Lazic SE (2008). Why we should use simpler models if the data allow this: relevance for ANOVA designs in experimental biology. BMC Physiology 8:16.

- Osman, A.G., F.I. Abd Elaziz and G.A. El Hassan, (2010). Effects of Biological and Mineral Fertilization on Yield, Chemical Composition and Physical Characteristics of Faba Bean (*Vicia faba* L.) Cultivar Seleim. Pakistan Journal of Nutrition, **9**(7): 703-708.
- Oweis,T; Hachum A., pala, M; (2004). lentil production under supplementary irrigation in Mediterranean environments .Agriculture waterManagement,Vol. **68**, 251-265
- Poth, M., J.S. Lafavre and D.D. Focht, (1986) Quantification by direct N¹⁵ dilution of fixed N₂ in incorporation into soil by pigeon pea (*Cajanus cajan*). Soil. Biol. Biochem., **189**: 125-127 .
- Salih, A.S.E., (1981). Development of new methods for preparing legume inoculant. M.Sc. (Agric), Thesis, University of Ain Shams, Cairo, Egypt
- Sanchez, P.A., (2002). Soil fertility and hunger in Africa. Sci, **295**: 2019-2020.
- Schultz, R.C., J.P. Colleti and R.R. Faltonson, (1998). Agroforestry opportunities for the United State of America. Agroforestry Systems. **31**: 117-142.
- Socalow, R.H., (1999). Nitrogen Management and the Future of Food. In: Lesson from the Management of Energy and Carbon. Proceedings of the National Academy of Sciences of the United States of America, **96**: 6001-6008.
- Vance, C.P., (2001). Symbiotic nitrogen fixation and phosphorus acquisitions. Plant nutrition in a world of declining renewable sources. Plant Physiol, **127**: 390-397.
- Watson, A.M., (1981). Faba bean production in the Sudan . A study of the economic and infrastructural context . ICARDA/IFAD Nile valley project, consultancy Report, pp: 103.
- Zhang,H.,pala, M.,Oweis,T.,Harris, H.C., (2000). Water use efficiency of chickpea and lentil in a Mediterranean environment Aust .J. Agric. Res. **51**.295-304.

